

On Some Perturbation Approaches to Population Dynamics

Francisco M. Fernández *
 INIFTA (UNLP, CCT La Plata-CONICET),
 División Química Teórica,
 Diag. 113 y 64 (S/N), Sucursal 4, Casilla de Correo 16,
 1900 La Plata, Argentina

June 3, 2008

Abstract

We show that the Adomian decomposition method, the time-series expansion, the homotopy-perturbation method, and the variational-iteration method completely fail to provide a reasonable description of the dynamics of the simplest prey-predator system.

It is well known that a reasonable description of the dynamics of a nonlinear system by means of a perturbation method is a difficult enterprise [1]. Recently, several approaches have been proposed for the treatment of the simplest model for the prey-predator interaction. They are based on the Adomian decomposition method [2], the time-series expansion [3], the homotopy-perturbation method [4, 5], and the variational-iteration method [6]. We have already shown that the time-power series given by the implementation of the homotopy-perturbation method proposed by Chowdhury et al [5] completely fails to yield the main features of the population dynamics [7]. In this short communication we briefly analyze the results of the other proposals in terms of what one expects from an approach designed to solve problems of population dynamics [1].

*e-mail: fernande@quimica.unlp.edu.ar

The chosen model is the prey–predator system [2–4, 6]

$$\begin{aligned}\frac{dx(t)}{dt} &= x(t)[a - by(t)], \quad a, b > 0 \\ \frac{dy(t)}{dt} &= -y(t)[c - dx(t)], \quad c, d > 0\end{aligned}\tag{1}$$

where $x(t)$ and $y(t)$ are the populations of rabbits and foxes, respectively. This nonlinear system exhibits a saddle point at $(x_s, y_s) = (0, 0)$ and a center at $(x_s, y_s) = (c/d, a/b)$ [1]. Besides, the populations obey the following curve in the $x - y$ plane:

$$\ln(x^c y^a) - dx - by = \ln(x_0^c y_0^a) - dx_0 - by_0\tag{2}$$

where x_0 and y_0 are the initial populations at time $t = 0$.

The homotopy–perturbation method proposed by Rafei et al [4] leads to time–series expansions of the form

$$\begin{aligned}x(t) &= x_0 + x_0(a - by_0)t + \dots \\ y(t) &= y_0 + y_0(dx_0 - c)t + \dots\end{aligned}\tag{3}$$

that are exactly the same as those obtained earlier by Biazar et al [3]. Obviously, they are suitable about the point (x_0, y_0) and will not predict the main features of the population dynamics (revealed at a much greater time span) which is what really matters in this field [1]. The variational–iteration method [6] yields more complicated expressions but the authors state that they agree with the time–power series mentioned above.

Fig. 1 shows the populations for Case I ($a = b = d = 1$, $c = 0.1$, $x_0 = 14$, $y_0 = 18$) in a time span larger than that considered in the earlier studies already mentioned above [2–4, 6]. We clearly appreciate that all those approaches predict a wrong behaviour of the populations. Besides, the time span in Fig. 1 is still not large enough to cover the whole portrait of the system evolution.

Case V ($a = b = c = d = 1$, $x_0 = 3$, $y_0 = 2$) is much more interesting [3, 4]. Fig. 2 shows the results in the $x - y$ plane. The time series [3, 4] predict a completely wrong behaviour and the resulting curve cross itself which can never happen as it is well known [1].

Summarizing: present analysis clearly shows that the Adomian decomposition method [2], the straightforward time–series expansion [3], the homotopy–

perturbation method [4,5], and the variational–iteration method [6] are completely useless for a reasonable prediction of the evolution of even the simplest prey–predator systems.

References

- [1] C. M. Bender and S. A. Orszag, Advanced mathematical methods for scientists and engineers, (McGraw-Hill, New York, 1978).
- [2] J. Biazar and R. Montazeri, Applied Mathematics and Computation 163 (2005) 841-847.
- [3] J. Biazar, M. Ilie, and A. Khoshkenar, Applied Mathematics and Computation 171 (2005) 486-491.
- [4] M. Rafei, H. Daniali, D. D. Ganji, and H. Pashaei, Appl. Math. Comput. 188 (2007) 1419-1425.
- [5] M. S. H. Chowdhury, I. Hashim, and O. Abdulaziz, Phys. Lett. A 368 (2007) 251-258.
- [6] E. Yusufoglu and B. Erbas, Phys. Lett. A 372 (2008) 3829-3835.
- [7] F. M. Fernández, Perturbation Theory for Population Dynamics, arXiv:0712.3376v1




Figure 1: Exact (solid, e) and series (dashed, s) populations for Case I




Figure 2: Exact (solid) and approximate (dashed) populations for Case V in the $x - y$ plane